

Communication between pharmacist and physician

Dr Guido Schmiemann PhD, MPH Department for Health Services Research Bremen University Germany



Current situation 30% of the german population take over the counter drugs Bundesgesundheitssurvey 2003 20 - 25% hospital admissions due to side effects Polypharmazie Report HKK 2012 One potentially serious drug interaction /day/ pharmacy Scharpf 2012 PZ

Current	situation
Country	Inhabitants/ pharmacy
Denmark	17.700
Netherlands	8.300
Slovenia	6.600
United Kingdom	4.600
Germany	3.900
Average EU	3.200
Belgium	2.100
ABDA-S ÖAK, AF	tatistik 2012; other countries- last year data according to ZAEU, SFK, IPU, IP, SPA
Dr Guido Schmiemann	

Current situation Germany

Pharmacists 48.422 (70.2% women)
Pharmacies 20.900

Physicians 138.472 (40% women)
GP 60.397 (39% women)

ABDA-Classifikationen


1. Contraindication 6 %
Serious side effects documented
Simvastatin - Erythromycin

2. Contraindication as a precaution 11%
Serious side effects theoreticaly possible
Levodopa- Metoclopramid

3. Monitoring or adjustment needed 46 %

4. In defined cases Monitoring or adjustment needed 4 %

ABDA-Datenbank (3/2012): 428 000 Interaktionen





Cross sectional study


" How do pharmacists and doctors judge their interprofessional communication regarding potentially serious interactions?


"How do they judge a standardized communication tool, developed to improve their communication? "

Dr Guido Schmieman









Physicians perspective		
	Disagree	Agree
"Pharmacies play in important role in the management of potentially serious interactions."	44,1% (n=41)	54,8% (n=51)
"Communication with other physicians is usually uneventful."	68,8% (n=64)	28,0% (n=26)
"Communication between pharmacy and medical practice is usually uneventful"	37,6% (n=35)	57,0% (n=53)
Dr Guido Schmiemann		

Pharmacists perspective	/e					
I	Never	2	3	4 5	Very	
	never	2	3		5	often
ask the patient to confer with his/her physician (before handing out the drug))	37,9%	34,5	6,9%	6,9%	6,9%	0 %
ask the patient to confer with his/her physician (after handing out the drug))	58,6%	20,7	3,4%	6,9%	6,9%	0 %
make a individual recommendation	31,0%	31,0%) 24,1%	0%	0%	3,4%
confer with physician while patient is waiting	3,4%	3,4%	13,8%	3,4%	34,5%	41,4%

Pharmacists "Dr not available- one can not pass the doctors assistant " "Interaction is known by the physician- not clinical relevant" Physicians Definition of serious interaction is relative" "Evaluation of interaction is incorrect" "They mostly contact us with bit and bops"

In wie weit stimmen Sie den folgenden Aussagen zu?	disagree	agree	missing
"A standardized communication tool (i.e template) can improve communication between pharmacies and physicians."	32/34%	65/56%	3/10%
"This tool can improve can improve communication between pharmacies and physicians ."	35/25%	65/67%	0/9%

Thank you schmiemann@uni-bremen.de

